

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.





**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**





Listing of Claims:






3. (Previously presented) A method for preparing a compound in an electronically excited state, comprising the steps of:





a) exposing a compound having the formula $D_1-\Pi-D_2$ to radiation, wherein D_1 and D_2 are electron donor groups; and Π comprises a bridge of π -conjugated bonds connecting D_1 and D_2 ; and

b) converting said compound to a multi-photon electronically excited state upon simultaneous absorption of at least two photons of said radiation by said compound, wherein the sum of the energies of all of said absorbed photons is greater than or equal to the transition energy from a ground state of said compound to said multi-photon excited state and wherein the energy of each absorbed photon is less than the transition energy between said ground state and the lowest single-photon excited state of said compound and is less than the transition energy between said multi-photon excited state and said ground state, wherein said compound is selected from the group consisting of










and mixtures thereof, where R=(CH₂)₁₁CH₃.

4. (Currently Amended) A method for preparing a compound in an electronically excited state, comprising the steps of:

- a) exposing a compound having the formula $D_1-\Pi-D_2$ to radiation, wherein D_1 and D_2 are electron donor groups; and Π comprises a bridge of π -conjugated bonds connecting D_1 and D_2 ; and
- b) converting said compound to a multi-photon electronically excited state upon simultaneous absorption of at least two photons of said radiation by said compound, wherein the sum of the energies of all of said absorbed photons is greater than or equal to the transition energy from a ground state of said compound to said multi-photon excited state and wherein the energy of each absorbed photon is less than the transition energy between said ground state and the lowest single-photon excited state of said compound and is less than the transition energy between said multi-photon excited state and said ground state, wherein said compound is further defined by a formula



where D_a is selected from the group consisting of N, O, S and P;

where D_b is selected from the group consisting of N, O, S and P;

m, n, o are integers such that $0 \leq m \leq 10$, $0 \leq n \leq 10$, $0 \leq o \leq 10$; and

where:

X, Y, Z are independently selected from the group consisting of: $CR_k=CR_l$; O; S; and $N-R_m$;

R_a, R_b, R_c, R_d are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{a1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl; where $0 < \alpha < 10$ and $1 < \beta < 25$, a group of aromatic rings having up to 20 carbons in the aromatic ring framework; fused aromatic rings, vinyl; allyl; 4-

styryl; acroyl; methacroyl; acrylonitrile, isocyanate; isothiocyanate; epoxides; strained ring olefins; $(-\text{CH}_2)_\delta\text{SiCl}_3$; $(-\text{CH}_2)_\delta\text{Si}(\text{OCH}_2\text{CH}_3)_3$; and $(-\text{CH}_2)_\delta\text{Si}(\text{OCH}_3)_3$; where $\delta < 25$;


Ra_1 , Ra_2 , and Ra_3 are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons, a functional group derived from an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof, ~~or~~ and methacryloyl chloride;

Re , Rf , Rg , Rh , Ri , Rj , Rk , Rl and Rm are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{OR}_{b1}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{NR}_{b2}\text{R}_{b3}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{CONR}_{b2}\text{R}_{b3}$, where R_{b1} , R_{b2} , and R_{b3} are independently selected from the group consisting of a **functional group derived from an amino acid; a polypeptide;** adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof ~~or~~ and methacryloyl chloride; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{CN}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{Cl}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{Br}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{I}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{-Phenyl}$, where $0 < \alpha < 10$ and $1 < \beta < 25$; a group of aromatic rings having up to 20 carbons in the aromatic framework; fused aromatic rings; CHO; CN; NO₂; Br; Cl; I; phenyl; an acceptor group containing more than two carbon atoms; a functional group derived from an amino acid and $\text{NR}_{e1}\text{R}_{e2}$; OR_{e3} ; where R_{e1} , R_{e2} , R_{e3} are defined as for R_n and R_o , where R_n and R_o are defined as any member of the group consisting of H; a linear or branched alkyl group with up to 25 carbons; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{OR}_{g1}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{NR}_{g2}\text{R}_{g3}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{CONR}_{g2}\text{R}_{g3}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{CN}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{Cl}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{Br}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{I}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{-Phenyl}$; aryl groups; fused aromatic rings; ~~polymerizable; and~~ polymerizable functionalities; functionalities; and

R_{g1} , R_{g2} , and R_{g3} are independently ~~selected from~~ selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; a functional group derived from an amino acid: a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof ~~or~~ and methacryloyl chloride.

5. (Currently Amended) A method for preparing a compound in an electronically excited state, comprising the steps of:

- a) exposing a compound having the formula $D_1-\Pi-D_2$ to radiation, wherein D_1 and D_2 are electron donor groups; and Π comprises a bridge of π -conjugated bonds connecting D_1 and D_2 ; and
- b) converting said compound to a multi-photon electronically excited state upon simultaneous absorption of at least two photons of said radiation by said compound, wherein the sum of the energies of all of said absorbed photons is greater than or equal to the transition energy from a ground state of said compound to said multi-photon excited state and wherein the energy of each absorbed photon is less than the transition energy between said ground state and the lowest single-photon excited state of said compound and is less than the transition energy between said multi-photon excited state and said ground state, wherein said compound is further defined by a formula



where D_a is selected from the group consisting of N, O, S and P;

where D_b is selected from the group consisting of N, O, S and P;

m, n, o are integers such that $0 \leq m \leq 10$, $0 \leq n \leq 10$, $0 \leq o \leq 10$; and

where:

X, Y, Z are independently selected from the group consisting of: $CR_k=CR_l$; O; S; and N- R_m ;

R_a , R_b , R_c , R_d are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{a1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl; where $0 < \alpha < 10$ and $1 < \beta < 25$, a group of aromatic rings having up to 20 carbons in the aromatic ring framework; fused aromatic rings, vinyl; allyl; 4-

styryl; acroyl; methacroyl; acrylonitrile, isocyanate; isothiocyanate; epoxides; strained ring olefins; $(-\text{CH}_2)_\delta\text{SiCl}_3$; $(-\text{CH}_2)_\delta\text{Si}(\text{OCH}_2\text{CH}_3)_3$; and $(-\text{CH}_2)_\delta\text{Si}(\text{OCH}_3)_3$; where $\delta < 25$;


R_{a1} , R_{a2} , and R_{a3} are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons, a functional group derived from an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof, ~~or~~ and methacryloyl chloride;

R_e , R_f , R_g , R_h , R_i , R_j , R_k , R_l and R_m are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{OR}_{b1}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{NR}_{b2}\text{R}_{b3}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{CONR}_{b2}\text{R}_{b3}$, where R_{b1} , R_{b2} , and R_{b3} are independently selected from a functional group derived from an amino acid, a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof ~~or~~ and methacryloyl chloride; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{CN}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{Cl}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{Br}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{I}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{-Phenyl}$, where $0 < \alpha < 10$ and $1 < \beta < 25$; a group of aromatic rings having up to 20 carbons in the aromatic framework; fused aromatic rings; CHO; CN; NO₂; Br; Cl; I; phenyl; an acceptor group containing more than two carbon atoms; a functional group derived from an amino acid and $\text{NR}_{e1}\text{R}_{e2}$; OR_{e3} ; where R_{e1} , R_{e2} , R_{e3} are defined as for R_n and R_o , where R_n and R_o are defined as any member of the group consisting of H; a linear or branched alkyl group with up to 25 carbons; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{OR}_{g1}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{NR}_{g2}\text{R}_{g3}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{CONR}_{g2}\text{R}_{g3}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{CN}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{Cl}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{Br}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{I}$; $(-\text{CH}_2\text{CH}_2\text{O})_\alpha(\text{CH}_2)_\beta\text{-Phenyl}$; aryl groups; fused aromatic rings; and polymerizable functionalities; functionalities; and

R_{g1} , R_{g2} , and R_{g3} are independently ~~selected from~~ selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; a functional group derived from an amino acid: a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof ~~or~~ and methacryloyl chloride.

6. (Currently Amended) A method for preparing a compound in an electronically excited state, comprising the steps of:

- a) exposing a compound having the formula $D_1-\Pi-D_2$ to radiation, wherein D_1 and D_2 are electron donor groups; and Π comprises a bridge of π -conjugated bonds connecting D_1 and D_2 ; and
- b) converting said compound to a multi-photon electronically excited state upon simultaneous absorption of at least two photons of said radiation by said compound, wherein the sum of the energies of all of said absorbed photons is greater than or equal to the transition energy from a ground state of said compound to said multi-photon excited state and wherein the energy of each absorbed photon is less than the transition energy between said ground state and the lowest single-photon excited state of said compound and is less than the transition energy between said multi-photon excited state and said ground state, wherein said compound is further defined by a formula



where D_a is selected from the group consisting of N, O, S and P;

where D_b is selected from the group consisting of N, O, S and P;

m, n, o are integers such that $0 \leq m \leq 10, 0 \leq n \leq 10, 0 \leq o \leq 10$; and

where:

X, Y, Z are independently selected from the group consisting of: $CR_k=CR_l$; O; S; and $N-R_m$;

R_a, R_b, R_c, R_d are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{a1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl; where $0 < \alpha < 10$ and $1 < \beta < 25$, a group of aromatic rings having up to 20 carbons in the aromatic ring framework; fused aromatic rings, vinyl; allyl; 4-styryl; acrolyl; methacroyl; acrylonitrile, isocyanate; isothiocyanate; epoxides; strained ring olefins; $(-CH_2)_\delta SiCl_3$; $(-CH_2)_\delta Si(OCH_2CH_3)_3$; and $(-CH_2)_\delta Si(OCH_3)_3$; where $\delta < 25$;

R_{a1} , R_{a2} , and R_{a3} are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons, a functional group derived from an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof, ~~or~~ and methacryloyl chloride;


R_e , R_f , R_i , R_j , R_k , R_l and R_m are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{b1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{b2}R_{b3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{b2}R_{b3}$, where R_{b1} , R_{b2} , and R_{b3} are independently selected from a functional group derived from an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof, ~~or~~, methacryloyl chloride; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl, where $0 < \forall \alpha < 10$ and $1 < \exists \beta < 25$; a group of aromatic rings having up to 20 carbons in the aromatic framework; fused aromatic rings; CHO; CN; NO₂; Br; Cl; I; phenyl; an acceptor group containing more than two carbon atoms; a functional group derived from an amino acid and $NR_{e1}R_{e2}$; OR_{e3} ; where R_{e1} , R_{e2} , R_{e3} are defined as for R_n and R_o , where R_n and R_o are defined as any member of the group consisting of H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{g1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{g2}R_{g3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{g2}R_{g3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl; aryl groups; fused aromatic rings; ~~polymerizable; and~~ polymerizable functionalities; functionalities; and

R_{g1} , R_{g2} , and R_{g3} are independently ~~selected from~~ selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; a functional group derived from an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof, ~~or~~ and methacryloyl chloride.

7. (Currently Amended) A method for preparing a compound in an electronically excited state, comprising the steps of:

a) exposing a compound having the formula $D_1-\Pi-D_2$ to radiation, wherein D_1 and D_2 are electron donor groups; and Π comprises a bridge of π -conjugated bonds connecting D_1 and D_2 ; and

b) converting said compound to a multi-photon electronically excited state upon simultaneous absorption of at least two photons of said radiation by said compound, wherein the sum of the energies of all of said absorbed photons is greater than or equal to the transition energy from a ground state of said compound to said multi-photon excited state and wherein the energy of each absorbed photon is less than the transition energy between said ground state and the lowest single-photon excited state of said compound and is less than the transition energy between said multi-photon excited state and said ground state, wherein said compound is further defined by a formula



where D_a is selected from the group consisting of N, O, S and P;

where D_b is selected from the group consisting of N, O, S and P;

m , n , o are integers such that $0 \leq m \leq 10$, $0 \leq n \leq 10$, $0 \leq o \leq 10$; and

where:

X , Y , Z are independently selected from the group consisting of: $CR_k=CR_l$; O; S; and N- R_m ;


R_a , R_b , R_c , R_d are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{a1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl; where $0 < \alpha < 10$ and $1 < \beta < 25$, a group of aromatic rings having up to 20 carbons in the aromatic ring framework; fused aromatic rings, vinyl; allyl; 4-styryl; acrolyl; methacroyl; acrylonitrile, isocyanate; isothiocyanate; epoxides; strained ring olefins; $(-CH_2)_\delta SiCl_3$; $(-CH_2)_\delta Si(OCH_2CH_3)_3$; and $(-CH_2)_\delta Si(OCH_3)_3$; where $\delta < 25$;

R_{a1} , R_{a2} , and R_{a3} are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons, a functional group derived from an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof, ~~or~~ and methacryloyl chloride;


R_e , R_f , R_i , R_j , R_k , R_l and R_m are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{b1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{b2}R_{b3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{b2}R_{b3}$, where R_{b1} , R_{b2} , and R_{b3} are independently selected from the group consisting of a functional group derived from an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof, ~~or~~ and methacryloyl chloride; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl, where $0 < \alpha < 10$ and $1 < \beta < 25$; a group of aromatic rings having up to 20 carbons in the aromatic framework; fused aromatic rings; CHO; CN; NO₂; Br; Cl; I; phenyl; an acceptor group containing more than two carbon atoms; a functional group derived from an amino acid and $NR_{e1}R_{e2}$; OR_{e3} ; where R_{e1} , R_{e2} , R_{e3} are defined as for R_n and R_o , where R_n and R_o are defined as any member of the group consisting of H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{g1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{g2}R_{g3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{g2}R_{g3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl; aryl groups; fused aromatic ring; ~~and polymerizable functionalities; functionalities; and~~
_____ R_{g1} , R_{g2} , and R_{g3} are independently ~~selected from~~ selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; a functional group derived from an amino acid: a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof, ~~or~~ and methacryloyl chloride.

13. (Currently Amended) A method for preparing a compound in an electronically excited state, comprising the steps of:


- a) exposing a compound having the formula $D_1-\Pi-D_2$ to radiation, wherein D_1 and D_2 are electron donor groups; and Π comprises a bridge of π -conjugated bonds connecting D_1 and D_2 ; and
- b) converting said compound to a multi-photon electronically excited state upon simultaneous absorption of at least two photons of said radiation by said compound, wherein the sum of the energies of all of said absorbed photons is greater than or equal to the transition energy from a ground state of said compound to said multi-photon excited state and wherein the energy of each absorbed photon is less than the transition energy between said ground state and the lowest single-photon excited state of said compound and is less than the transition energy between said multi-photon excited state and said ground state, wherein said compound is further defined by a formula




where A_a and A_b ~~can be~~ are independently selected from: selected from the group consisting of: CHO; CN; NO₂, and




A1




A2




A3




A4




A5




A6




A7




A8




A9




A10




A11




A12




A13




A14




A15




A16




A17




A18




A19




A20




A21




A22




A23




A24




A25




A26




A27




A28




A29




A30




A31




A32




A33




A34




A35




A36




A37




A38




A39



A40



A41



A42

in addition A_a and A_b can be independently selected from the group consisting of: Br, Cl, and I; and where $0 \leq m \leq 10$, $0 \leq n \leq 10$, $0 \leq o \leq 10$; and where:

X, Y, Z are independently selected from the group consisting of: $\text{CR}_k=\text{CR}_l$; O; S; and N-R_m ;

R_a , R_b , R_c , R_d are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{OR}_{a1}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{NR}_{a2}\text{R}_{a3}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{CONR}_{a2}\text{R}_{a3}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{CN}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{Cl}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{Br}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{I}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{-Phenyl}$; where $0 < \alpha < 10$ and $1 < \beta < 25$, a group of aromatic rings having up to 20 carbons in the aromatic ring framework; fused aromatic rings, vinyl; allyl; 4-styryl; acroyl; methacryl; acrylonitrile, isocyanate; isothiocyanate; epoxides; strained ring olefins; $(-\text{CH}_2)_\delta\text{SiCl}_3$; $(-\text{CH}_2)_\delta\text{Si}(\text{OCH}_2\text{CH}_3)_3$; and $(-\text{CH}_2)_\delta\text{Si}(\text{OCH}_3)_3$; where $\delta < 25$;

R_{a1} , R_{a2} , and R_{a3} are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons, a functional group derived from an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof, ~~or~~ and methacryloyl chloride;

R_e , R_f , R_i , R_j , R_k , R_l and R_m are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{OR}_{b1}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{NR}_{b2}\text{R}_{b3}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{CONR}_{b2}\text{R}_{b3}$, where R_{b1} , R_{b2} , and R_{b3} are independently selected from the group consisting of a functional group derived from an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof, ~~or~~ methacryloyl chloride; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{CN}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{Cl}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{Br}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{I}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{-Phenyl}$, where $0 < \forall \alpha < 10$ and $1 < \exists \beta < 25$; a group of aromatic rings having up to 20 carbons in the aromatic framework; fused aromatic rings; CHO; CN; NO₂; Br; Cl; I; phenyl; an acceptor group containing more than two carbon atoms; a functional group derived from an amino acid and $\text{NR}_{e1}\text{R}_{e2}$; OR_{e3} ; where R_{e1} , R_{e2} , R_{e3} are defined as for R_n and R_o , where R_n and R_o are defined as any member of the group consisting of H; a linear or branched alkyl group with up to 25 carbons; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{OR}_{g1}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{NR}_{g2}\text{R}_{g3}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{CONR}_{g2}\text{R}_{g3}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{CN}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{Cl}$;


$-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{Br}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{I}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{-Phenyl}$; aryl groups; fused aromatic rings; and polymerizable functionalities; functionalities; and

R_{g1} , R_{g2} , and R_{g3} are independently ~~selected from:~~selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; a functional group derived from an amino acid; or a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof ~~or~~ and methacryloyl chloride.


14. (Currently Amended) A method for preparing a compound in an electronically excited state, comprising the steps of:

a) exposing a compound having the formula $\text{D}_1\text{-}\Pi\text{-D}_2$ to radiation, wherein D_1 and D_2 are electron donor groups; and Π comprises a bridge of π -conjugated bonds connecting D_1 and D_2 ; and


b) converting said compound to a multi-photon electronically excited state upon simultaneous absorption of at least two photons of said radiation by said compound, wherein the sum of the energies of all of said absorbed photons is greater than or equal to the transition energy from a ground state of said compound to said multi-photon excited state and wherein the energy of each absorbed photon is less than the transition energy between said ground state and the lowest single-photon excited state of said compound and is less than the transition energy between said multi-photon excited state and said ground state, wherein said compound is further defined by a formula




where A_a and A_b ~~can be~~are independently ~~selected from:~~selected from the group consisting of: CHO; CN; NO_2 , and




A1




A2




A3




A4




A5




A6




A7




A8




A9




A10




A11




A12




A13




A14




A15




A16




A17




A18




A19




A20




A21




A22




A23




A24




A25




A26




A27




A28




A29




A30




A31




A32




A33




A34




A35




A36




A37




A38




A39



A40



A41



A42

in addition A_a and A_b can be independently selected from the group consisting of: Br, Cl, and I; and where $0 \leq m \leq 10$, $0 \leq n \leq 10$, $0 \leq o \leq 10$; and where:

X, Y, Z are independently selected from the group consisting of: $CR_k=CR_l$; O; S; and $N-R_m$;

R_a , R_b , R_c , R_d are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{a1}$;
 $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{a2}R_{a3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$;
 $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$;
 $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl; where $0 < \alpha < 10$ and $1 < \beta < 25$, a group of aromatic rings having up to 20 carbons in the aromatic ring framework; fused aromatic rings, vinyl; allyl; 4-styryl; acroyl; methacryl; acrylonitrile, isocyanate; isothiocyanate; epoxides; strained ring olefins; $-(CH_2)_\delta SiCl_3$; $-(CH_2)_\delta Si(OCH_2CH_3)_3$; and $-(CH_2)_\delta Si(OCH_3)_3$; where $\delta < 25$;

R_{a1} , R_{a2} , and R_{a3} are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons, a functional group derived from an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof, ~~or~~ and methacryloyl chloride;

R_e , R_f , R_i , R_j , R_k , R_l and R_m are independently selected from the group consisting of: H; a linear or branched alkyl group with up to 25 carbons; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{b1}$;
 $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{b2}R_{b3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{b2}R_{b3}$, where R_{b1} , R_{b2} , and R_{b3} are independently selected from the group consisting of a functional group derived from an amino acid; a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof, ~~or~~ and methacryloyl chloride;
 $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Br$;
 $-(CH_2CH_2O)_\alpha-(CH_2)_\beta I$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta$ -Phenyl, where $0 < \forall \alpha < 10$ and $1 < \exists \beta < 25$; a group of aromatic rings having up to 20 carbons in the aromatic framework; fused aromatic rings; CHO; CN; NO₂; Br; Cl; I; phenyl; an acceptor group containing more than two carbon atoms; a functional group derived from an amino acid and $NR_{e1}R_{e2}$; OR_{e3} ; where R_{e1} , R_{e2} , R_{e3} are defined as for R_n and R_o , where R_n and R_o are defined as any member of the group consisting of H; a linear or branched alkyl group with up to 25 carbons;
 $-(CH_2CH_2O)_\alpha-(CH_2)_\beta OR_{g1}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta NR_{g2}R_{g3}$;
 $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CONR_{g2}R_{g3}$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta CN$; $-(CH_2CH_2O)_\alpha-(CH_2)_\beta Cl$;

$-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{Br}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{I}$; $-(\text{CH}_2\text{CH}_2\text{O})_\alpha-(\text{CH}_2)_\beta\text{-Phenyl}$; aryl groups; fused aromatic rings; ~~polymerizable; and polymerizable functionalities; functionalities; and~~

$\text{R}_{\text{g}1}$, $\text{R}_{\text{g}2}$, and $\text{R}_{\text{g}3}$ are independently ~~selected from:~~selected from the group consisting of:
H; a linear or branched alkyl group with up to 25 carbons; a functional group derived from an amino acid; or a polypeptide; adenine; guanine; tyrosine; cytosine; uracil; biotin; ferrocene, ruthenocene, cyanuric chloride and derivatives thereof ~~or~~ and methacryloyl chloride.